

EXCELLENCE

F1 Hybrid Sweet Pepper

OUTSTANDING QUALITIES

- ◆ COMPACT PLANTS WITH GOOD LEAF COVER
- ◆ LARGE BLOCKY FRUIT
- ◆ HIGH RESISTANCE TO TMV, CMV AND PVY
- ◆ IDEAL FOR OPEN FIELD PRODUCTION


Excellence is a widely adapted, early, hybrid blocky pepper with a compact growth habit and good leaf cover. The fruit are large, strongly four lobed, blocky, show very little purpling and ripen to a uniform glossy red. Fruit set is concentrated, resulting in high first pick yields. **Excellence** has high resistance to Tobacco mosaic (TMV), Cucumber mosaic (CMV) and Potato Y (PVY). **Excellence** is recommended for open field production, but can also be trellised using the "Spanish" method under protection.



SPECIAL VARIETAL REQUIREMENTS

- As fruit set is concentrated, we suggest that flowers of the 1st and 2nd bottom splits are removed to allow the plant to maintain a good balance between leaf cover, fruit load and roots

CHARACTERISTIC*	EXCELLENCE
KIND	F1 hybrid pepper (<i>Capsicum L.</i>)
TYPE	Blocky, California Wonder type
MATURITY	Early
FRUIT DIMENSIONS	Approximately 10 x 10 cm (app. 250 g)
FRUIT SHAPE	Blocky
FRUIT WALL	Thick
SMOOTHNESS	Smooth surface with medium deep lobes, shoulders and blossom end
FRUIT COLOUR	Dark green turning glossy red
PLANT TYPE	Compact
DISEASE REACTION (SCIENTIFIC)	High resistance: <i>Tobacco mosaic virus (TMV)</i> ; <i>Cucumber mosaic virus (CMV)</i> and <i>Potato Y virus (PVY)</i>
PRODUCTION	Under protection (shade net) and open field
POPULATION GUIDE	Open field: 20 000 – 35 000 plants per ha Under protection: 2 – 2.3 stems per m ²
USE	Bulk packaging
SPECIAL FEATURES	Uniformly blocky, large fruit and ideal for open field production without trellising

* Characteristics given are affected by production methods such as soil type, nutrition, planting population, planting date and climatic conditions. Please read disclaimer.
 WARNING: VARIETY PROTECTED UNDER **PLANT BREEDERS RIGHTS**. UNAUTHORIZED MULTIPLICATION AND/OR MARKETING OF SEED PROHIBITED.

Disclaimer: This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Sakata Seed Southern Africa (Pty) Ltd Conditions of Sale before ordering seed.

Resistance: is the ability of a plant variety to restrict the growth and development of a specified pest or pathogen and/or the damage they cause when compared to susceptible plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure (HR = High resistance, IR = Intermediate resistance).

Recent version: Kindly contact Sakata or Area Representative for the most recent version of this Technical Bulletin.

GENERAL TIPS FOR SWEET PEPPER PRODUCTION

Seedling production

- The Sakata Range Test is available for hybrid peppers and is used to determine suitable germination temperatures of a specific seed lot.
- Cold sensitivity: low temperatures have a detrimental effect on germination. Seed will not germinate at temperatures below 15 °C.
- Pepper seedlings should be transplanted before root growth becomes stunted. It is suggested to transplant seedlings from 200-cavity trays when the first pair of true leaves is fully extended. Seedlings in larger cavities can be transplanted later.
- Make sure to plant seedlings in deep enough cavities to avoid J-roots and subsequent poor growth.

Poor shelf life

- Wilted fruit has a compromised shelf life, so avoid harvesting in the heat of the day. Cold storage conditions should be maintained at a consistent 8.3 – 9 °C.
- Temperature fluctuations lead to condensation and bacterial rotting, higher temperatures lead to ripening, and lower temperatures result in cold injury.

Flat fruit

Causes

- The occurrence of short-blocky or flat fruit is common for some varieties under high temperature conditions
- Low temperatures can cause flat fruit as fertilisation does not take place when temperature is too low
- Incidence of short blocky fruit increases with high N-concentration in the nutrient solution

Control

- Select varieties suited to the environmental conditions.

Fruit cracking

- This defect is often a problem with high diurnal temperature swings and high humidity at night, for example in spring and autumn.
- Fruit become very turgid at night as a result of water uptake. If the humidity is very high, little transpiration takes place and eventually fruit cracks as a result of the pressure in the fruit wall.
- Reduce humidity in the greenhouse through ventilation before sunrise.
- Control greenhouse temperature to 18 – 30 °C, or produce during a cooler time of year.

- Keep the ratio ppm N-NO₃ : ppm N-NH₄ to around 5:1 and limit the N-NH₄ concentration to < 32 ppm.

Climatic requirements

- Peppers grow best when relative humidity (RH) is 65 – 75 %
- Maintain good ventilation (0.5 m/s) to keep conditions favourable for transpiration
- Pepper plants need good light (1 100 – 1 300 µs/m² or 60 000 lux). Heavy shade can induce stress, but light shade stimulates growth
- The ideal temperature is around 18 °C (minimum) and 25 °C (maximum)
- Temperatures lower than 15 °C result in very poor growth
- Temperatures higher than 28 °C induce stress

Despite the need for warm conditions the plant is sensitive to high temperatures. Above 32 °C the flowers are inclined to fall off and few fruits, if any, set at temperatures above 35 °C, especially when these high temperatures are coupled with dry winds. Fruit that form at such high temperatures is usually malformed. The fruit is also very sensitive to sunburn and for this reason Sakata has select varieties that develop well leaf covering to protect the fruit.

Unmarketable fruit

Malformed, sun scalded, cracked or diseased fruits must be removed as soon as possible to allow the plant to set new fruit. If plants lose leaves due to disease, young fruit should be removed so the plant can restore the balance between fruit (nutrient sink) and leaves (nutrient source).

- Do not irrigate in the late afternoon.
- Application of Calcium as a foliar spray every two weeks can reduce cracking.

Sun scald

Sudden exposure of fruit to high light intensity (mainly the UV spectrum) can cause sun scald (sunken, pale tissue that often becomes infected by secondary pathogens). This is more of a problem in open field production, and when foliage cover is reduced. Avoid over pruning of the canopy and use varieties with adequate foliage cover.

Disclaimer: This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Sakata Seed Southern Africa (Pty) Ltd Conditions of Sale before ordering seed.

Resistance: is the ability of a plant variety to restrict the growth and development of a specified pest or pathogen and/or the damage they cause when compared to susceptible plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure (HR = High resistance, IR = Intermediate resistance).

Recent version: Kindly contact Sakata or Area Representative for the most recent version of this Technical Bulletin.