


# Sowing Guideline: Cauliflower & Broccoli

## Eastern Cape

VARIETY	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
<b>CAULIFLOWER</b>												
Eskimo												
Incline												
<b>BROCCOLI</b>												
Arcadia												
Imagine												
Marathon												
Parthenon												
Pharos												

 Suggested sowing time: Sowing time applies to the area of seedling nursery and farm location

### Factors causing physiological disorders in cauliflower & broccoli

- Wrong sowing time
- Cold temperatures, especially below 7°C
- Excessive fertilisation of seedlings
- Cold grown seedlings
- Oversized seedlings at transplant
- Difference in temperatures between seedling nursery and farm
- Growth checks caused by heat, drought, water logging and disease
- Diurnal temperature swings

Disclaimer: This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Sakata Seed Southern Africa (Pty) Ltd Conditions of Sale before ordering seed.

Recent version: Kindly contact Sakata or Area Representative for the most recent version of this Sowing Guide.

Ref: Caul Broc Sow South Cape: 31-07-14