

ICICLE Radish

OUTSTANDING QUALITIES

- STRONG TALL TOPS
- **EXCEPTIONAL UNIFORM ROOTS**
- WIDELY ADAPTED
- **GOOD YIELD POTENTIAL**
- MILD FLAVOUR

Icicle is a blunt cylindrical, pure white radish with a mild flavour. Icicle is adapted to a wide range of soils and geographical regions. The relative maturity is 25 - 30 days after sowing. Icicle has very strong tops and light green in





SPECIAL VARIETAL REQUIREMENTS

- Suitable for all year round production
- Contact the area representative for more information

CHARACTERISTIC*	ICICLE
TYPE	White radish (Raphanus sativus L. var. sativus L.)
MATURITY	Medium (around 25 - 30 days from sowing)
SEASON	Year round production
ROOT SHAPE	Cylindrical
DIMENSIONS	2.5 x 10 - 15 cm
SKIN COLOUR	Pure white
SMOOTHNESS	Very smooth
FLESH COLOUR	Pure white
FLESH TEXTURE	Very firm, crisp
FLAVOUR	Mild
TOP	Tall strong foliage
TOP HEIGHT	15 - 25 cm
TOP COLOUR	Light green
DISEASE REACTION (SCIENTIFIC)	
FIELD HOLDING	Very good
YIELD POTENTIAL	Very good
SUGGESTED POPULATION	10 – 12 kg per ha, 1.2 - 1.8 million seeds per ha
MARKET USE	Bunching, salad packs and pre-pack
SPECIAL FEATURES	High quality roots with exceptional uniformity, mild flavour

^{*} Characteristics given are affected by production methods such as soil type, nutrition, planting population, planting date and climatic conditions. Please read disclaimer.

Disclaimer: This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Sakata Seed Southern Africa (Pty) Ltd Conditions of Sale before ordering seed.

Resistance: is the ability of a plant variety to restrict the growth and development of a specified pest or pathogen and/or the damage they cause when compared to susceptible plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure (HR = High resistance, IR

Recent version: Kindly contact Sakata or Area Representative for the most recent version of this Technical Bulletin.

Tel. +27 11 548 2800

Copyright: (not to be reproduced) Fax. +27 11 548 2820

Email. info.saf@sakata.eu Website. www.sakata.com

